ACTIVE INGREDIENT (per ml):

Lidocaine Hydrochloride 20m

Purilled water

Inactives: Methylparaben,

Propylparaben, Sodium Chloride, Citric Acid

WARNINGS

For external use only. Avoid contact with eyes. If allergic reaction occurs, discontinue use and consult a physician.

Do not use in large quantities particularly over raw or blistered areas. Indicated for use on firs degree burns only. Consult a physician if burn related pain persists for more than 7 days, or i symptoms improve and reoccur within a few days If swallowed, get medical help or contact a poisor control center right away.

MANUFACTURED BY

Radix Laboratories, Inc 1334 International Drive Eau Claire, WI 54701

1 900 027 2240



NDC: 50480-037-25



FAST ACTING

BURN RELIEF

TOPICAL ANALGESIC





Keep away from children

NET CONTENTS 8.5 fl. oz (250 ml)

PRECAUTIONS

Avoid prolonged storage at cold temperatures, which can cause preservatives to precipitate. To clarify, place product in warm water bath (100°F - 120°F) for about 30 minutes.

DIRECTIONS

For temporary, first-aid relief of pain associated with minor scalds and bums. Adults and childrer 2 years and older; apply to affected area no more than 3-4 times daily. Consult a physician before administering to children under 2 years Can be sprayed or poured on.

STORAGE:

Store at room temperature between 59°F and 86°F (15°C - 30°C)